

# The “Spot Sign”: A predictor of hematoma expansion

Mckenzie Merritt DO<sup>1</sup>, Trusha Mehta DO<sup>1</sup>, Jasmine Chopra DO<sup>1</sup>, Adena Shahinian DO<sup>1</sup>,

Aroucha Vickers DO<sup>1,2</sup>, Robert Balsiger DO<sup>1,2</sup>, Paul Janda DO, JD<sup>1,2</sup>

Valley Hospital Medical Center - Las Vegas, NV<sup>1</sup>, Las Vegas Neurology Center<sup>2</sup>

## INTRODUCTION

In this poster, we show the utility of the CT Angiogram (CTA) “Spot sign” as a predictor of intracranial hematoma expansion and subsequent neurologic decline, and demonstrate the measures we utilized to decrease the morbidity and mortality of the patient in this case report. We discuss the hospital course of a 69-year-old male who presented with hypertensive emergency and focal neurologic deficits and was found to have a large basal ganglia hypertensive bleed with “Spot sign” present on CTA head and neck with and without contrast. Patient was treated very aggressively and was given hypertonic solutions due to the team’s quick recognition of the “Spot sign” signaling potential for future hematoma expansion. Though as predicted by the “Spot sign”, his bleed did expand, the patient survived with our aggressive measures at the onset of bleed.

## BACKGROUND


The “Spot sign” is a radiographic sign that can be found on a CTA study and represents a spot of active extraluminal extravasation and accumulation of contrast dye within an intracerebral hemorrhage (1, 2). Theoretically, the sign has been seen in shape as linear, branching or spherical (3). When this sign is identified on imaging, it indicates an increased likelihood of significant increase in size of hematoma expansion, as this sign is an independent predictor of growth and poor outcomes (4). Meta-analysis studies have shown that, "for each 10% increase in hematoma volume, patients were 5% more likely to die and 16% more likely to increase one point on the modified rankin scale" (5). However, if identified correctly the “Spot sign” can be utilized to identify patients who are most likely to benefit from medical and surgical therapy for rapid expansion of intracranial hematoma (6, 7). The “Spot sign” serves as a predictor for possible hematoma expansion and warrants close neurologic monitoring.

## REFERENCES

1. Wada R, Aviv RI, Fox AJ, et al. CT angiography “spot sign” predicts hematoma expansion in acute intracerebral hemorrhage. *Stroke*. 2007 Apr;38(4):1257-62. doi: 10.1161/01.STR.0000259633.59404.f3. PMID: 17322083
2. Demchuk AM, Dowlatahahi D, Rodriguez-Luna D, et al. Prediction of haematoma growth and outcome in patients with intracerebral haemorrhage using the CT-angiography spot sign (PREDICT): a prospective observational study. *Lancet Neurol*. 2012 Apr;11(4):307-14. doi: 10.1016/S1474-4422(12)70038-8. PMID: 22405630
3. CT angiographic spot sign (intracerebral hemorrhage). Website. <https://radiopaedia.org/articles/ct-angiographic-spot-sign-intracerebral-haemorrhage?lang=us>. Published 2021. Accessed March 20, 2021.
4. Soun JE, Montes D, Yu F, et al. Spot Sign in Secondary Intraventricular Hemorrhage Predicts Early Neurological Decline. *Clin Neuroradiol*. 2019 Nov 27. doi: 10.1007/s00062-019-00857-2. Epub ahead of print. PMID: 31776595.
5. Spontaneous intracerebral hemorrhage: Treatment and prognosis. Web site. [https://www.uptodate.com/contents/spontaneous-intracerebral-hemorrhage-treatment-and-prognosis?search=spot%20sign%20on%20CTA&source=search\\_result&selectedTitle=2~150&usage\\_type=default&display\\_rank=2](https://www.uptodate.com/contents/spontaneous-intracerebral-hemorrhage-treatment-and-prognosis?search=spot%20sign%20on%20CTA&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2). Published 2021. Accessed March 20, 2021.
6. Ciura VA, Brouwers HB, Pizzolato R, et al. Spot sign on 90-second delayed computed tomography angiography improves sensitivity for hematoma expansion and mortality: prospective study. *Stroke*. 2014 Nov;45(11):3293-7. doi: 10.1161/STROKEAHA.114.005570. PMID: 25300974; PMCID: PMC4213346
7. Brouwers HB, Battery TW, Musial HH, et al. Rate of Contrast Extravasation on Computed Tomographic Angiography Predicts Hematoma Expansion and Mortality in Primary Intracerebral Hemorrhage. *Stroke*. 2015 Sep;46(9):2498-503. doi: 10.1161/STROKEAHA.115.009659. PMID: 26243220; PMCID: PMC4550492
8. Du FZ, Jiang R, Gu M, et al. The accuracy of spot sign in predicting hematoma expansion after intracerebral hemorrhage: a systematic review and meta-analysis. *PLoS One*. 2014 Dec 26;9(12):e115777. doi: 10.1371/journal.pone.0115777. PMID: 25541717; PMCID: PMC4277365

## CASE PRESENTATION

This is a 68-year-old right-handed Caucasian male with no known prior medical history other than alcohol abuse who presented secondary to a sudden onset of left sided weakness while sitting on a barstool. Patient fell to the ground, did not lose consciousness, and came in for potential large vessel occlusion. On the initial stat CT head without contrast, the patient was noted to have a large right supra-tentorial 6.4 x 3.1 cm hemorrhage (Image A). On arrival, the patient was noted to be hypertensive with systolic blood pressure as high as 195. A CTA head and neck with and without contrast was ordered which demonstrated that he did not have evidence of vascular malformation but did have evidence of large blush of contrast within the hematoma consistent with active hemorrhage (Image B). At this time the patient was started on hypertonic solutions in the form of 3% hypertonic saline in an urgent fashion. Repeat CT head demonstrated a mild interval increase in the size of the right basal ganglia hemorrhage with intraventricular extension (Image C). With serial CT head imaging without contrast, the hemorrhage remained stable, and the hypertonic saline was eventually discontinued. His hospital course was complicated by delirium tremens in the setting of daily alcoholism, for which he was also treated for and subsequently required intubation and eventual peg and trach placement. Further neurologic workup including EEGs were unremarkable for epileptiform discharges. He continued to remain stable from a neurological standpoint. Patient was able to be discharged from the hospital to a long-term acute care facility with clear instructions for repeat CT head imaging to be done as an outpatient.



**Image A.** Initial CT head without contrast demonstrating right basal ganglia hemorrhage

**Image B.** The initial CTA head and neck with contrast demonstrating large blush of contrast within the hematoma

**Image C.** Repeat CT head without contrast demonstrating worsening of right basal ganglia hematoma with intraventricular extension

## DISCUSSION/CONCLUSION

The CTA “Spot sign” in intracranial hemorrhage has been shown to be a reliable imaging sign for determining hematoma expansion (8). This sign is an important predictor for pending neurologic decline secondary to hemorrhagic expansion and is an indicator for patients who may have improved outcomes with aggressive and immediate interventions to avoid significant morbidity and mortality (4). One such intervention is for craniotomy with hematoma removal which can lead to an improved prognosis in “Spot sign” positive patients (9).

It is important to recognize patients with intracranial hemorrhage and have CTA performed as soon as possible as it has been shown that the longer it takes for the patient to get the CTA, the more the positive predictive value of the “Spot sign” decreases (10). In fact, CTA followed by 90 second delayed post-contrast CT, also known as a “delayed CTA”, is a further predictor of the amount of hematoma expansion based on the amount of contrast extravasation, showing that the “Spot sign” can measure active bleeding in these patients which can help better select patients who may see a benefit from additional therapies (6, 7). There is also importance to the timing of the CTA, as when the “Spot sign” was seen arterial and therefore earlier in the imaging, there was greater expansion of the hemorrhage while the later or venous images improved the detection of “Spot signs” (11).

We present the importance of recognizing the “Spot sign” to help alert clinicians to the utility of this important diagnostic tool. This is a clear sign of impending hematoma expansion and prompt recognition is required to guide prognosis and management to decrease the morbidity and mortality otherwise expected from a rapidly expanding bleed. In our presented case, the quick actions of the team in recognizing the “Spot sign” and beginning 3% hypertonic saline early led to only mild interval increase in the size of the right basal ganglia hemorrhage with some intraventricular extension as opposed to the catastrophic possibilities that the “Spot sign” signifies.

**Acknowledgements:** We would like to thank our patient who allowed us to care for him and report his case. We would also like to thank the entire team that was involved in the care of our patient.

## REFERENCES

9. Ge C, Zhao W, Guo H, et al. Comparison of the clinical efficacy of craniotomy and craniopuncture therapy for the early stage of moderate volume spontaneous intracerebral haemorrhage in basal ganglia: Using the CTA spot sign as an entry criterion. *Clin Neurol Neurosurg*. 2018 Jun;169:41-48. doi: 10.1016/j.clineuro.2018.04.002. Epub 2018 Apr 3. PMID: 29625339.
10. Dowlatahahi D, Brouwers HB, Demchuk AM, et al. Predicting Intracerebral Hemorrhage Growth With the Spot Sign: The Effect of Onset-to-Scan Time. *Stroke*. 2016 Mar;47(3):695-700. doi: 10.1161/STROKEAHA.115.012012. PMID: 26846857; PMCID: PMC4766058
11. Rodriguez-Luna D, Dowlatahahi D, Aviv RI, et al. Venous phase of computed tomography angiography increases spot sign detection, but intracerebral hemorrhage expansion is greater in spot signs detected in arterial phase. *Stroke*. 2014 Mar;45(3):734-9. doi: 10.1161/STROKEAHA.113.003007. PMID: 24481974